

TOX

PRE CARE

No clot reducing meds 10 – 14 days before

Risks/Complications Allergic reaction, abnormal facial expressions, paralysis of nearby muscles, eyelid and eye brow droop, disorientation, double vision, facial pain, asymmetrical appearance, head ache, nausea, flu-like symptoms, impaired swallowing or speech, permanent muscle atrophy, nerve irritability, and production of antibodies with unknown general health effects. **Contraindications** Myasthenia Gravis, MS, ALS, Guillen- Barre, any other neurological or movement disorder, hypersensitivity to Type A Toxin, or its constituents human albumin, use of aminoglycoside antibiotics, or spectinomycin, pregnancy or breast feeding.

TOX

POST CARE



- Lying down for 6 hours
- Massage appointments within 24 hrs
- Pressure on area for the day
- Makeup for one hour
- Strenuous exercise
- Hair washing, blow drying, or shower for the day
- Air travel day of procedure

Onset 2 – 3 days, max effect 10 –14 days. Duration approximately 3 months. The period between each treatment should be at least 3 months. Seek medical help immediately if swallowing, speech, or respiratory issues occur. Injection-related reactions occur and may include: bruising, swelling, tenderness, and headache. Increased bruising and bleeding are common in patients taking medications that reduce clotting.